# **Elabscience**®

## Recombinant Human Interleukin-2/IL-2 (C-6His-Avi) Biotinylated

### Catalog Number: PKSH033973

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Human	
Source	HEK293 Cells-derived Human Interleukin-2; IL-2 protein Ala21-Thr153, with an C-	
	terminal His & Avi	
Calculated MW	18.3 kDa	
Observed MW	16-18 kDa	
Accession	P60568	
Bio-activity	Measured in a cell proliferation assay using CTLL- 2 mouse cytotoxic T cells. The E	
	$_{50}$ for this effect is 0.2 ng/ml.	
Properties		
Purity	> 90 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

#### Data

kDa	МК	R
120 90	-	
60		
40	-	
30	-	
20	-	
14	-	

> 90 % as determined by reducing SDS-PAGE.

#### Background

Interleukin-2(IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system, belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.