

A Reliable Research Partner in Life Science and Medicine

Recombinant Rat sCD14 Protein(Fc Tag)

Catalog Number: PDMR100052

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Rat **Species**

Source HEK293 Cells-derived Rat CD14 protein Ser18-Tyr341, with an C-terminal FC

Calculated MW 60.5 kDa Observed MW 60-80 kDa Accession Q63691

Not validated for activity **Bio-activity**

Properties

> 90% as determined by reducing SDS-PAGE. **Purity**

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -Storage

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of


reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% **Formulation**

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Rat CD14 proteins, 2 µg/lane of Recombinant Rat CD14 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 60.5KD

Background

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules associating with an the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 14 (CD14) is a member of the CD system. It takes its name from its inclusion in the CD molecule surface marker proteins. CD14 exists in two forms: a form anchored into the membrane or a soluble form. CD14 was found expressed in macrophages, neutrophil granulocyte and dendritic cells. The major function is to serve as a co-receptor (along with an TLR4 and MD-2) for the bacterial lipopolysaccharide (LPS) and other pathogen-associated molecular patterns.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com