Elabscience®

CREB1 Polyclonal Antibody

catalog number: E-AB-31024

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Reactivity	Human;Mouse;Rat
Immunogen	Synthesized peptide derived from human CREB-1 around the non-phosphorylation
minunogen	site of Ser133.
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein
Duitt	protectant and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
Data	PC-3 Hela 3T3
Data	
Data 138- MCF7 100- 55- 40- 35- 25- 15-	PC-3 Hela 3T3
Data 170- 100- 55- 40- 35 25 15 Western Blot analy	PC-3 Hela 3T3
Data MCF7 100- 55- 40- 35- 25- 15- Western Blot analy Polyclonal An	PC-3 Hela 3T3 CREB-1 creB-1 sis of various cells using CREB1
Data MCF7 TO- TO- TO- TO- TO- TO- TO- TO-	PC3 Hela 3T3 CREB-1 creB-1 sis of various cells using CREB1 tibody at dilution of 1:1000.
Data MCF7 TO- TO- TO- TO- TO- TO- TO- TO-	PC3 Hela 313 CREB-1 creB-1 creB-1 creB-1 tibody at dilution of 1:1000. rved-MW:37 kDa Hated-MW:37 kDa
Data MGF7 100- 55- 40- 35- 55- 15- Western Blot analy Polyclonal An Obse Calcu	PC3 Hela 313 CREB-1 creB-1 creB-1 creB-1 tibody at dilution of 1:1000. rved-MW:37 kDa Hated-MW:37 kDa
Data MCF7 TO- TO- TO- TO- TO- TO- TO- TO-	PC3 Hela 3T3 CREB-1 CREB-1 cred-MW:37 kDa Hated-MW:37 kDa

Background

This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Implicated in synchronization of circadian rhythmicity.