

Recombinant Mouse Peroxiredoxin 5/PRDX5 Protein (His Tag)

Catalog Number: PKSM040619

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	E.coli-derived Mouse Peroxiredoxin 5/PRDX5 protein Met 49-Leu 210, with an N-terminal His
Calculated MW	18.5 kDa
Observed MW	18.5 kDa
Accession	P99029-1
Bio-activity	Measured by its ability to reduce H ₂ O ₂ . The specific activity is > 100 pmoles/min/μg.

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 92 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.5

Peroxiredoxin-5, also known as Alu corepressor 1, Antioxidant enzyme B166, Liver tissue 2D-page spot 71B, Peroxisomal antioxidant enzyme, Thioredoxin peroxidase PMP20, Thioredoxin reductase, PRDX5 and ACR1, is cytoplasm protein that belongs to the peroxiredoxin 2 family. Peroxiredoxin-5 / PRDX5 reduces hydrogen peroxide and alkyl hydroperoxides with reducing equivalents provided through the thioredoxin system. Peroxiredoxin-5 / PRDX5 is involved in intracellular redox signaling. The Peroxiredoxins / Prx are a family of 25 kDa peroxidases that can reduce H₂O₂ using an electron from thioredoxin (Trx) or other substances. The mammalian Peroxiredoxins / Prx family is divided into six groups (PRDX1, PRDX2, PRDX3, PRDX4, PRDX5, PRDX6) on the basis of homology of amino acid sequences. They are located in the cytosol and play a role in the cell signaling system. All six mammalian peroxiredoxins are expressed in the lung. Peroxiredoxins / Prx is overexpressed in breast cancer tissues to a great extent suggesting that Peroxiredoxins / Prx has a proliferative effect and may be related to cancer development or progression.