

Recombinant Human Rac2 protein (His Tag)

Catalog Number: PDEH100997

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human Rac2 protein Met1-Cys189, with an N-terminal His & C-terminal His
Calculated MW	20.7 kDa
Observed MW	25 kDa
Accession	P15153
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ras-related C3 botulinum toxin substrate 2 (Rac2) is a small G-protein belonging to the Ras subfamily of the GTPase family. Rac2 acts as an "on / off" switch for signal transduction cascades and motilities. When GDP is attached to the small G-protein, the enzyme is inactivated. Release of the GDP and replace of the GTP activate the GTPase. Rac2 remains active until the GTP is hydrolyzed to GDP. Rac2 is a hematopoietic-specific Rho family GTPase implicated as an important constituent of the NADPH oxidase complex and shares 92% amino acid identity with the ubiquitously expressed Rac1. The small G-protein Rac2 regulates the rearrangements of actin and membrane necessary for Fcγ receptor-mediated phagocytosis by macrophages. Activated Rac2 binds to the p21-binding domain of PAK1 and this binding provided a basis for microscopic methods to localize activation of these G proteins inside cells.

For Research Use Only