

## WWOX Polyclonal Antibody

catalog number: E-AB-65948

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |                                                                                    |
|---------------------|------------------------------------------------------------------------------------|
| <b>Reactivity</b>   | Human;Mouse;Rat                                                                    |
| <b>Immunogen</b>    | Recombinant fusion protein of human WWOX (NP_057457.1).                            |
| <b>Host</b>         | Rabbit                                                                             |
| <b>Isotype</b>      | Rabbit IgG                                                                         |
| <b>Purification</b> | Affinity purification                                                              |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications

### Recommended Dilution

|           |               |
|-----------|---------------|
| <b>WB</b> | 1:1000-1:2000 |
|-----------|---------------|

### Data



Western blot analysis of extracts of various cell lines using WWOX Polyclonal Antibody at dilution of 1:3000.

**Observed-MW:46 kDa**

**Calculated-MW:4 kDa/21 kDa/23 kDa/26 kDa/35 kDa/41 kDa/46 kDa**

### Preparation & Storage

|                 |                                                                                                        |
|-----------------|--------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                                        |
| <b>Shipping</b> | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

### Background

This gene encodes a member of the short-chain dehydrogenases/reductases (SDR) protein family. This gene spans the FRA16D common chromosomal fragile site and appears to function as a tumor suppressor gene. Expression of the encoded protein is able to induce apoptosis, while defects in this gene are associated with multiple types of cancer. Disruption of this gene is also associated with autosomal recessive spinocerebellar ataxia 12. Disruption of a similar gene in mouse results in impaired steroidogenesis, additionally suggesting a metabolic function for the protein. Alternative splicing results in multiple transcript variants.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.8