

Purified Anti-Mouse CD3ε Antibody[145-2C11]

Catalog Number: GF1103A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD3ε protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone	145-2C11
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 µg/mL (0.5×10 ⁶ -1×10 ⁶ cells)
-----	--

Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD3ε Antibody[145-2C11](Right) and 0.2 µg Armenian Hamster IgG, κ Isotype Control(Left),

followed by Elab Fluor® 647-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody, then anti-Mouse CD8a FITC-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD3ε is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3ε forms a TCR complex by associating with the CD3δ, γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.