A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL-2RB/CD122 Protein (His Tag)

Catalog Number: PKSH032571

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| \mathbf{r} | | crip | | | | | |
|--------------|--------------|------|----|---|----|---|---|
| | 00 | 0 | РΤ | n | ŤΤ | | m |
| v | \mathbf{c} | v. | | w | w | w | ш |

Species Human

Source HEK293 Cells-derived Human IL-2RB;CD122 protein Ala27-Asp239, with an C-terminal

His

Calculated MW 25.6 kDa
Observed MW 35-40 kDa
Accession P14784

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Human IL-2RB; also known as interleukin-2 receptor subunit beta; is the receptor for interleukin-2. IL2 receptor complex is involved in receptor mediated endocytosis and transduces the mitogenic signals of IL2. IL2 receptor complex has three forms with respect to ability to bind IL2. IL-2RB is belonged to a type I membrane protein; and has a 26 residue signal peptide; a 214 residue extracellular region; a 25 residue transmembrane region and a 286 residue cytoplasmic domain. IL-2RB is the subunit critical for receptor-mediated signaling via physically or functionally coupling to other signaling molecules; such as the Jak-STAT and Src-family protein tyrosine kinase although it lacks apparent catalytic motifs.

For Research Use Only