

Recombinant Human FGF-2/FGFb Protein (aa 132-288)

Catalog Number: PKSH032437

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human FGF-2;FGFb protein Gly132-Ser288
Calculated MW	17.4 kDa
Observed MW	16 kDa
Accession	P09038-4
Bio-activity	Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED ₅₀ for this effect is 0.42 ng/ml.


Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.01 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, 3% Trehalose, 4% Mannitol, pH 7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.



Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED₅₀ for this effect is 0.42 ng/ml.

Background

FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation; growth; and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis; cell growth; pattern formation; embryonic development; metabolic regulation; cell migration; neurotrophic effects; and tissue repair. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain; pituitary; kidney; retina; bone; testis; adrenal gland liver; monocytes; epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b; 1c; 2c; 3c and 4.

For Research Use Only