A Reliable Research Partner in Life Science and Medicine

## **Recombinant Casein Kinase 2 beta Monoclonal Antibody**

catalog number: AN301820L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human;Rat;Mouse

Immunogen Recombinant human Casein Kinase 2 beta fragment

Host Rabbit Isotype lgG, κ Clone A532


**Purification** Protein Apurified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

#### **Applications Recommended Dilution**


1:500-1:1000 **WB IHC** 1:50-1:100 1:50 IF


#### Data



Western Blot with Casein Kinase 2 beta Monoclonal Immunohistochemistry of paraffin-embedded Human colon Antibody at dilution of 1:1000. Lane 1: HeLa, Lane 2: MCF- cancer using Casein Kinase 2 beta Monoclonal Antibody at 7, Lane 3: NIH/3T3, Lane 4: C6 dilution of 1:100.

> Observed-MW:25 kDa Calculated-MW:25 kDa





Immunohistochemistry of paraffin-embedded Rat cerebrum using Casein Kinase 2 beta Monoclonal Antibody at dilution Immunofluorescent analysis of (100% Ice-cold methanol) fixed NIH/3T3 cells using anti-Casein Kinase 2 beta Monoclonal Antibody at dilution of 1:50.

Rev. V1.1

#### **Preparation & Storage**

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** Ice bag

#### **Background**

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Casein Kinase 2 beta is a ubiquitous protein kinase which regulates metabolic pathways, signal transduction, transcription, translation, and replication. The enzyme is composed of three subunits, alpha, alpha prime and beta, which form a tetrameric holoenzyme. The alpha and alpha prime subunits are catalytic, while the beta subunit serves regulatory functions. The enzyme localizes to the endoplasmic reticulum and the Golgi apparatus. It participates in Wnt signaling, and plays a complex role in regulating the basal catalytic activity of the alpha subunit.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.1