



Recombinant TFPI/LACI/EPI Monoclonal Antibody

catalog number: AN300196P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human TFPI Protein
Host	Rabbit
Isotype	IgG
Clone	B114
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS
Applications	Recommended Dilution
WB:	1:1000; ICC/IF: 1:100; IP: 1:20-1:50

Data	
 Western blot analysis of extracts from HepG2 cells at 1:1000. Observed-MW:40-55 kDa Calculated-MW:35 kDa	 Immunofluorescence analysis of HepG2 cells labelling TFPI. The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (shown in green). Nuclear DNA was labeled in blue with DAPI. Control: Secondary antibody only.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background
This gene encodes a Kunitz-type serine protease inhibitor that regulates the tissue factor (TF)-dependent pathway of blood coagulation. The coagulation process initiates with the formation of a factor VIIa-TF complex, which proteolytically activates additional proteases (factors IX and X) and ultimately leads to the formation of a fibrin clot. The product of this gene inhibits the activated factor X and VIIa-TF proteases in an autoregulatory loop. Inhibition of the encoded protein restores hemostasis in animal models of hemophilia. This gene encodes multiple protein isoforms that differ in their inhibitory activity, specificity and cellular localization. [provided by RefSeq, Jul 2016]

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.4