# **Elabscience**®

## **DNM2** Polyclonal Antibody


### catalog number: E-AB-65692

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human DNM2
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution


WB	1:500-1:2000
IF	1:50-1:200

#### Data




Western blot analysis of extracts of various cell lines using DNM2 Polyclonal Antibody at 1:500 dilution.

### Observed-MW:110 kDa Calculated-MW:97 kDa/98 kDa



Immunofluorescence analysis of U2OS cells using DNM2 Polyclonal Antibody at dilution of 1:150. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using DNM2 Polyclonal antibody at dilution of 1:150. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of PC-12 cells using DNM2 Polyclonal antibody at dilution of 1:150. Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

#### Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

# **Elabscience**®

Dynamins represent one of the subfamilies of GTP-binding proteins. These proteins share considerable sequence similarity over the N-terminal portion of the molecule, which contains the GTPase domain. Dynamins are associated with microtubules. They have been implicated in cell processes such as endocytosis and cell motility, and in alterations of the membrane that accompany certain activities such as bone resorption by osteoclasts. Dynamins bind many proteins that bind actin and other cytoskeletal proteins. Dynamins can also self-assemble, a process that stimulates GTPase activity. Five alternatively spliced transcripts encoding different proteins have been described. Additional alternatively spliced transcripts may exist, but their full-length nature has not been determined.

For Research Use Only