

APC Anti-Mouse CD170 Antibody[S17007L]

Catalog Number: AN00629E

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|-------------------------|--|
| Reactivity | Mouse |
| Host | Rat |
| Isotype | Rat IgG1, κ |
| Clone No. | S17007L |
| Isotype Control | APC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822E] |
| Conjugation | APC |
| Conjugation Information | APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |

Applications

| Applications | Recommended usage |
|--------------|--|
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |

Data



Staining of C57BL/6 murine bone marrow cells with FITC Anti-Mouse/Human CD11b Antibody and APC Anti-Mouse CD170 Antibody[S17007L] (left) or APC Rat IgG1, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

| | |
|----------|---|
| Storage | Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |

Antigen Information

| | |
|-----------------|----------------------|
| Alternate Names | OB BP2;CD33L2;OB-BP2 |
| Uniprot ID | Q920G3 |
| Gene ID | 8778 |

For Research Use Only

Background

CD170, also known as Siglec-F, Siglec-5, is a member of the Sialic acid-binding Ig-like lectin family, type I single pass transmembrane protein, with 4 extracellular Ig-like domains and 2 ITIM motifs in the cytoplasmic domain; preferentially binds [alpha]-2,3-linked sialic acid. Siglec F is expressed in eosinophils, alveolar macrophages and intestinal microfold (M) cells and induces apoptosis of the lung eosinophils during allergic asthma.

For Research Use Only