

## Recombinant Rat CD83/HB15 Protein (Fc Tag)

**Catalog Number:** PKSR030225

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat CD83/HB15 protein Met1-Ala134, with an C-terminal hFc
<b>Calculated MW</b>	39.4 kDa
<b>Observed MW</b>	53 kDa
<b>Accession</b>	B2GV95
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 76 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 76 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD83 is considered as a marker of mature dendritic cells as well as an adhesion receptor that binds to resting monocytes and a subset of activated CD8+ T cells. In certain conditions, CD83 tended to dimerize or even multimerize through its aberrant intermolecular disulfide bonds. The injection of CD83-Ig can significantly enhance the rate of tumor growth and inhibit the T cell growth.