

A Reliable Research Partner in Life Science and Medicine

PE Anti-Rat CD71 Antibody[OX-26]

Catalog Number: AN00622D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat
Host Mouse

Isotype Mouse IgG2a, κ

Clone No. OX-26

Isotype Control PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of SD Rat bone marrow cells with PE Anti-Rat CD71 Antibody[OX-26] (filled gray histogram) or PE Mouse IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Transferrin Receptor;T9

Uniprot ID Q99376

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

7037

CD71 is a 95 kD type II transmembrane protein. It is expressed on proliferating cells, reticulocytes, and brain endothelium. It is involved in the activation, proliferation, and iron metabolism of cells.

Fax: 1-832-243-6017