Elabscience®

Human MMP-8 Antibody Pair Set

Catalog No.E-KAB-0055ApplicationsSynonymsMMP8, CLG1, HNC, PMNL-CL, Neutrophil Collagenase

ELISA

Kit components & Storage

Title	Specifications	Storage
Human MMP-8 Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year.
		Avoid freeze / thaw cycles.
Human MMP-8 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0055)		
		Human MMP-8 Capture Antibody	Human MMP-8 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human MMP-8 protein	Recombinant Human MMP-8 protein	
Information	Swissprot	P22894		
Product details	Reactivity	Human	Human	
	Host	Rabbit	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A	Protein A	
	Specificity	Detects Human MMP-8 in ELISAs.		

For Research Use Only

Elabscience®

Applications

Human MMP-8 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human MMP-8 Capture Antibody	
Capture			
ELISA	1:1000-1:10000	Human MMP-8 Detection	Optical Density
Detection		Antibody (Biotin)	
			0.01 0.1 1 10 100 Human MMP-8 concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the enzyme encoded by this gene is stored in secondary granules within neutrophils and is activated by autolytic cleavage. Its function is degradation of type I, II and III collagens. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.