A Reliable Research Partner in Life Science and Medicine

Recombinant Human E-Cadherin/CDH1 Protein (His Tag)

Catalog Number: PKSH033437

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip					
	00	CI	411	n	П	n	m
JU	\mathbf{c}	U		J A	ш	v.	ш

Species Human

Source HEK293 Cells-derived Human E-Cadherin/CDH1 protein Asp155-Ile707, with an C-

terminal His

Calculated MW61.2 kDaObserved MW80-90 kDaAccessionP12830

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

E-Cadherin is a classical member of the cadherin superfamily. The encoded protein is a calcium-dependent cell-cell adhesion glycoprotein composed of five extracellular cadherin repeats; a transmembrane region; and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric; breast; colorectal; thyroid; and ovarian cancers. Loss of function is thought to contribute to progression in cancer by increasing proliferation; invasion; and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells; and the cytoplasmic domain is required for internalization. Identified transcript variants arise from mutation at consensus splice sites. Also; E-Cadherin has a potent invasive suppressor role and it is a ligand for integrin alpha-E/beta-7.

For Research Use Only