



A Reliable Research Partner in Life Science and Medicine

# **Mouse TIMP-1 Antibody Pair Set**

Catalog No. E-KAB-0287 Applications ELISA

**Synonyms** TIMP1, CLGI, EPA, EPO, HCI

## **Kit components & Storage**

Title	Specifications	Storage
Mouse TIMP-1 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse TIMP-1 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0287)		
		Mouse TIMP-1 Capture Antibody	Mouse TIMP-1 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Mouse TIMP-1 protein	Recombinant Mouse TIMP-1 protein	
Information	Swissprot	P12032		
Product details	Reactivity	Mouse	Mouse	
	Host	Rat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Mouse TIMP-1 in ELISAs.	·	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

### **Applications**

Mouse TIMP-1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse TIMP-1 Capture Antibody	
Capture			Alisa
ELISA Detection	1:1000-1:10000	Mouse TIMP-1 Detection Antibody (Biotin)	0.01 000 10000 100000 Mouse TIMP-1 concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### **Background**

This gene belongs to the TIMP gene family. The proteins encoded by this gene family are natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is transcribed in the opposite direction.