

PE/Cyanine7 Anti-Human CD99 Antibody[HI156]

Catalog Number: E-AB-F1339H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone No.	HI156
Isotype Control	PE/Cyanine7 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802H]
Conjugation	PE/Cyanine 7
Conjugation Information	PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm (e.g., a 780/60 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



Human peripheral blood lymphocytes are stained with PE/Cyanine7 Anti-Human CD99 Antibody[HI156] (filled gray histogram) or PE/Cyanine7 Mouse IgG2a, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	E2 antigenHBA71MIC2MSK5X;CD99
Uniprot ID	P14209
Gene ID	4267

For Research Use Only

Background

CD99 is a type I single chain transmembrane protein devoid of N-linked glycosylation sites encoded by the pseudoautosomal gene MIC2. CD99 has an apparent molecular weight of 32 kD and is widely expressed on a variety of tissues. CD99 is highly expressed on thymocytes, T cells, and T cell leukemias and lymphomas. However, it is absent on some B cell lines, fetal B cells, eosinophils, granulocytes and the NK-cell line YT. CD99 is involved in spontaneous rosette formation with erythrocytes and may also be involved in other T-cell and hematopoietic cell adhesion pathways. CD99 has been reported to activate a caspase-independent death pathway in T cells under some conditions. CD99 interacts with a number of proteins including ferritin heavy chain 1, karyopherin beta 1, TRIP13, cyclophilin A, annexin II, and ubiquitin-conjugating enzyme E2H.