

## Recombinant Human STAT6 Protein (Baculovirus, His Tag)

Catalog Number: PKSH030708

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Baculovirus-Insect Cells-derived Human STAT6 protein Met 1-Trp 847, with an C-terminal His
<b>Calculated MW</b>	95.5 kDa
<b>Observed MW</b>	100 kDa
<b>Accession</b>	P42226-1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 20% glycerol, 0.3mM DTT. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

Signal transducer and activator of transcription 6 (STAT6) is a transcription factor that is activated by interleukin-4 (IL-4)-induced tyrosine phosphorylation and mediates most of the IL-4-induced gene expression. STAT6 plays a central role in exerting interleukin-4 (IL-4) mediated biological responses and is found to induce the expression of BCL2L1/BCL-XL, which is responsible for the anti-apoptotic activity of IL4. Transcriptional activation by STAT6 requires the interaction with coactivators like p300 and the CREB-binding protein (CBP). NF-κB and tyrosine-phosphorylated Stat6 can directly bind each other in vitro and in vivo, which suggests that the direct interaction between Stat6 and NF-κB may provide a basis for synergistic activation of transcription by IL-4 and activators of NF-κB.

### For Research Use Only