

Recombinant Phospho-HSP27 (Ser78) Monoclonal Antibody

catalog number: AN302087L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen phosphorylated human HSP27 (Ser78) peptide

 Host
 Rabbit

 Isotype
 IgG, κ


 Clone
 A811

Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Applications	Recommended Dilution
WB	1:500-1:1000
IHC	1:100-1:500
IF	1:50
IP	1:50-1:100


Data




Western Blot with Phospho-HSP27 (Ser78) Monoclonal Immunohistochemistry of paraffin-embedded Human breast Antibody at dilution of 1:1000. Lane 1: HeLa + Anisomycin(- cancer using Phospho-HSP27 (Ser78) Monoclonal Antibody), 2 HeLa+ Anisomycin(+) at dilution of 1:500.

Observed-MW:27 kDa Calculated-MW:23 kDa






Immunohistochemistry of paraffin-embedded Human cervical cancer using Phospho-HSP27 (Ser78) Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human tonsil using Phospho-HSP27 (Ser78) Monoclonal Antibody at dilution of 1:500.



For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: w w w .elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Immunofluorescent analysis of (100% Ice-cold methanol) fixed A: HeLa, B HeLa+ Anisomycin(+) cells using anti-Phospho-HSP27 (Ser78) Monoclonal Antibody at dilution of

Immunoprecipitation analysis using anti-Phospho-HSP27 (Ser78) Monoclonal Antibody. Western blot was performed from the immunoprecipitate using Phospho-HSP27 (Ser78) Monoclonal Antibody at a dilution of 1:100. Lane 1: 5% Input, Lane 2: Phospho-HSP27 Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG Isotype

Observed-MW:27 kDa Calculated-MW:23 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Hsp27 is one of the small HSPs that are constitutively expressed at different levels in various cell types and tissues. Like other small HSPs, HSP27 is regulated at both the transcriptional and posttranslational levels. In response to stress, the HSP27 expression increases several-fold to confer cellular resistance to the adverse environmental change. HSP27 is phosphorylated at Ser15, Ser78, and Ser82 by MAPKAPK-2 as a result of the activation of the p38 MAP kinase pathway. Phosphorylation of HSP27 causes a change in its tertiary structure, which shifts from large homotypic multimers to dimers and monomers. It has been shown that phosphorylation and increased concentration of HSP27 modulates actin polymerization and reorganization.

For Research Use Only