



A Reliable Research Partner in Life Science and Medicine

## PerCP/Cyanine5.5 Anti-Human CD172a/b Antibody[SE5A5]

Catalog Number: AN00317J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_		4.0	
	escri	MTI.	on.
ш	COUL		UII

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. SE5A5

PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792J] Isotype Control

PerCP/Cyanine 5.5 Conjugation

**Conjugation Information** PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected

using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

## **Applications** Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM** 

> of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

## **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

## **Antigen Information**

**Alternate Names** SIRPa;SIRPb;SIRPalpha/beta;BIT;SHPS1;MFR;P84;PTPNS1;CD172 antigen-like family

member A;CD172 antigen-like family member B;

**Uniprot ID** O00241 Gene ID 140885

**Background** CD172a, also known as signal-regulatory protein α (SIRPα), src homology 2 domain-

> containing phosphatase substrate-1 (SHPS1), PTPNS1, BIT, MFR, and P84, is a 75-110 kD transmembrane glycoprotein involved in receptor tyrosine kinase coupled signaling pathway. It belongs to the lg superfamily and is primarily expressed on monocytes/macrophages, granulocytes, dendritic cells, and neurons. CD172a serves as a substrate of activated receptor tyrosine kinases (RTKs). The interaction of CD172a intracellular domain with SHP-1 and SHP-2 displays negative signaling in the

regulation of leukocyte adhesion and transmigration, T cell activation, macrophage fusion, and phagocytosis. CD47 (IAP) is the extracellular ligand for CD172a. SIRPa was recently demonstrated to be a specifc marker for cardiomyocytes derived from

human pluripotent stem cells2.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Rev. V1.6