

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Violet 540 Anti-Human CD16 Antibody[3G8]

Catalog Number: E-AB-F1236T3

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Rhesus; Cynomolgus

Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. 3G8

Isotype Control Elab Fluor® Violet 540 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-

F09792T31

Conjugation Elab Fluor® Violet 540

**Conjugation Information** Elab Fluor® Violet 540 is designed to be excited by the violet laser (405 nm) and detected

using an optical filter centered near 548 nm (e.g., a 572/28 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


## Applications Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### **Data**



Staining of normal human peripheral blood cells with APC

Anti-Human CD56/NCAM Antibody[5.1H11] and Elab Fluor® Violet 540 Anti-Human CD16 Antibody[3G8](left) or Elab

Fluor<sup>®</sup> Violet 540 Mouse IgG1, κ Isotype Control(right). Cells in the lymphocytes gate were used for analysis.

# **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

## **Antigen Information**

Alternate Names Fc gamma receptor;Fc gamma receptor 3;FcyRIII

**Uniprot ID** P08637;O75015;

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

941

Elabscience®

CD16 is a 60 kD highly glycosylated protein. It is a member of the lg superfamily and is also known as B7-1, B7, and Ly-53. CD16 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD16 and CD86 (B7-2) on antigen presenting cells ( such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD16 appears to be expressed later in the immune response than CD86. CD16 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.