

## Recombinant Human APOM Protein (His Tag)

**Catalog Number:** PKSH033762

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                      |   |
|----------------------|---|
| <b>Species</b>       | Human   |
| <b>Source</b>        | HEK293 Cells-derived Human APOM protein Met1-Asn188, with an C-terminal His |
| <b>Calculated MW</b> | 22.3 kDa  |
| <b>Observed MW</b>   | 26 kDa  |
| <b>Accession</b>     | O95445  |
| <b>Bio-activity</b>  | Not validated for activity  |

### Properties

|                       |  |
|-----------------------|--|
| <b>Purity</b>         | > 90 % as determined by reducing SDS-PAGE.   |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.  |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.  |
|                       | Please refer to the specific buffer information in the printed manual.   |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.   |

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

Apolipoprotein M is a secreted protein which belongs to the Lipocalin family. ApoM often presents in high density lipoprotein (HDL) and to a lesser extent in triglyceride-rich lipoproteins (TGRLP) and low density lipoproteins (LDL). The ApoM gene encoded protein is expressed in liver and kidney; secreted through the plasma membrane but remains membrane-bound. ApoM probably involved in lipid transport. ApoM can bind sphingosine-1-phosphate; myristic acid; palmitic acid and stearic acid; retinol; all-trans-retinoic acid and 9-cis-retinoic acid. The expression of ApoM could be regulated by platelet activating factor (PAF); Transforming Growth Factors (TGF); Insulin-Like Growth factor (IGF) and Leptin.

### For Research Use Only