# Purified Anti-Human CD340 Antibody[SER4]

catalog number: AN003530P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

**Reactivity** Human

Immunogen Recombinant Human CD340 protein

**Host** Mouse

**Isotype** Mouse IgG2b, κ

Clone SER4

**Purification** >98%, Protein A/G purified

**Conjugation** Unconjugated


buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze

to completely remove the stabilizer prior to labeling.

## Applications Recommended Dilution

FCM  $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$ 

#### Data



MCF-7 were stained with 0.2 μg Purified Anti-Human CD340 Antibody[SER4] (Right) and 0.2 μg Mouse IgG2b, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

#### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

#### Background

ErbB2, also called Neu and Her2, is a transmembrane glycoprotein in the ErbB family of tyrosine kinase receptors for EGF superfamily growth factors. ErbB2 is widely expressed in epithelial cells and over-expressed in a large number of breast carcinomas. ErbB2 has no identified ligands but heterodimerizes with ErbB1/EGF R, ErbB3, or ErbB4 to form higher affinity signaling complexes. The protease ADAM10 releases a 110 kDa soluble fragment of ErbB2 from the cell surface. ErbB2 plays roles in development, cancer, communication at the neuromuscular junction, and regulation of cell growth and differentiation. The ErbB2/ErbB3 heterodimer is expressed in the majority of breast, skin, ovary and gastrointestinal tumors and transduces a highly mitogenic signal in response to neuregulin 1 (NRG1; heuregulin 1) or NRG2. ErbB3, ErbB2 and neuregulin are all required for formation of the sympathetic nervous system.

### For Research Use Only