

## Recombinant Human GASP-1/WFIKKNRP Protein(His Tag)

**Catalog Number:** PDMH100312

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Mammalian-derived Human GASP-1/WFIKKNRP proteins Leu35-His576,with an C-terminal His
<b>Calculated MW</b>	59.5 kDa
<b>Observed MW</b>	80 kDa
<b>Accession</b>	Q8TEU8
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human GASP-1/WFIKKNRP proteins , 2µg/lane of Recombinant Human GASP-1/WFIKKNRP proteins was resolved with SDS-PAGE under reducing conditions , showing bands at 80 KD

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.7

WAP, kazal, immunoglobulin, kunitz and NTR domain-containing protein 2, also known as Growth and differentiation factor-associated serum protein 1, WAP, follistatin, immunoglobulin, kunitz and NTR domain-containing-related protein, WFIKKN-related protein, WFIKKN2 and GASPI, is a secreted protein that belongs to the WFIKKN family. WFIKKN2 contains two BPTI/Kunitz inhibitor domains, one Ig-like C2-type (immunoglobulin-like) domain, one Kazal-like domain, one NTR domain and one WAP domain. WFIKKN2 is primarily expressed in ovary, testis and brain, but not in liver. In fetal tissues, it is primarily expressed in brain, skeletal muscle, thymus and kidney. WFIKKN2 is a protease-inhibitor that contains multiple distinct protease inhibitor domains. It probably has serine protease- and metalloprotease-inhibitor activity. It inhibits the biological activity of mature myostatin, but not activin. WFIKKN2 protein binds mature GDF8/myostatin and myostatin propeptide and inhibits the biological activity of myostatin.