Recombinant Aconitase 1/ACO1 Monoclonal Antibody

catalog number: **AN301871L**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant human Aconitase 1/ACO1 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C131493
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications	Recommended Dilution
WB	1:500-1:2000


Data



Western Blot with Aconitase 1/ACO1 Monoclonal Antibody at dilution of 1:2000. Lane 1: Mouse kidney, Lane 2: Rat kidney

Observed-MW:105 kDa

Calculated-MW:98 kDa



Western Blot with Aconitase 1/ACO1 Monoclonal Antibody at dilution of 1:2000. Lane 1: Human liver tissue lysate, the Aconitase1/ACO1 Rabbit mAb pre-adsorbed with 3μM of the synthetic peptides, Lane 2: Human liver tissue lysate, the Aconitase 1/ACO1 Rabbit mAb with no peptide blocking.

Observed-MW:105 kDa

Calculated-MW:98 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Aconitase 1 is a bifunctional, cytosolic protein that functions as an essential enzyme in the TCA cycle and interacts with mRNA to control the levels of iron inside cells. When cellular iron levels are high, this protein binds to a 4Fe-4S cluster and functions as an aconitase. Aconitases are iron-sulfur proteins that function to catalyze the conversion of citrate to isocitrate. When cellular iron levels are low, the protein binds to iron-responsive elements (IREs), which are stem-loop structures found in the 5′ UTR of ferritin mRNA, and in the 3′ UTR of transferrin receptor mRNA.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com