



A Reliable Research Partner in Life Science and Medicine

# Purified Anti-Human CD11a Antibody[TS1/22.1.1.13]

catalog number: E-AB-F1055A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Reactivity Human

Immunogen Recombinant Human CD11a protein

**Host** Mouse

 $\begin{tabular}{lll} \textbf{Isotype} & Mouse IgG1, \kappa \\ \textbf{Clone} & TS1/22.1.1.13 \end{tabular}$ 

**Purification** >98%, Protein A/G purified


**Buffer** Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

# Applications Recommended Dilution

**FCM** 2  $\mu$ g/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

#### Data



Human peripheral blood Granulocytes were stained with 0.2  $\mu$ g Purified Anti-Human CD11a Antibody[TS1/22.1.1.13] (Right) and 0.2  $\mu$ g Mouse IgG1,  $\kappa$  Isotype Control(Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

# **Preparation & Storage**

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

## **Background**

CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1 $\alpha$  chain and integrin  $\alpha$ L subunit. CD11a non-covalently associates with integrin  $\beta$ 2 (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

For Research Use Only