

A Reliable Research Partner in Life Science and Medicine

Recombinant Human cytomegalovirus (HCMV) Glycoprotein B / gB Protein (His Tag)

Catalog Number: PKSV030208

Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						٠.		
D	e	8	C	۲T	n	tı	M I	ī

Species CMV

Source HEK293 Cells-derived CMV cytomegalovirus (HCMV) Glycoprotein B / gB protein

Arg 777-Val 907, with an C-terminal His

 Mol_Mass
 93.0 kDa

 Accession
 AAA45920.1

Bio-activity Not validated for activity

Properties

Purity > 70 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 70 % as determined by reducing SDS-PAGE.

Background

Cytomegalovirus (CMV) (human herpesvirus 5) glycoprotein B, also referred as CMV gB or gB, which belongs to the herpesviridae glycoprotein B family. It is a 97-amino acid glycoprotein encoded by the ORF of UL55. Cytomegalovirus Glycoprotein B protein is the most abundant component of the envelope, a target of neutralizing antibodies with at least two defined neutralizing epitopes and an essential replication component. Cytomegalovirus Glycoprotein B protein plays important roles in HCMV entry, cell-cell spread of internal virions, and fusion of infected cells. In addition, Cytomegalovirus Glycoprotein B protein is one envelope protein capable of heparin binding. It forms a physical association with host cell annexin II independent of the presence of calcium.

For Research Use Only

Fax: 1-832-243-6017