

PE Anti-Mouse CD36 Antibody[HM36]

Catalog Number: E-AB-F1291D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone No.	HM36
Isotype Control	PE Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	--

Data



C57BL/6 murine abdominal macrophages elicited by starch broth are stained with PE Anti-Mouse CD36 Antibody[HM36] (filled gray histogram) or PE Armenian Hamster IgG Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	FAT;gpIIIb;gpIV
Uniprot ID	Q08857
Gene ID	12491

For Research Use Only

Background

CD36 is a 85 kD glycoprotein, also known as FAT, gpIIIb, or gpIV. It is a member of the class B scavenger receptor family, expressed on platelets, monocytes, macrophages, megakaryocytes, microvasculature, dendritic cells and mammary endothelial cells. The primary ligands for CD36 have been reported to be oxidized low density lipoprotein, anionic phospholipids, and collagens I, IV, and V. CD36 acts as a scavenger receptor thus promoting the removal of apoptotic neutrophils and other apoptotic bodies, as well as clearance of defective erythrocytes.