

## Recombinant Human CD274 Protein(His Tag)

**Catalog Number:** PDMH100256

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Mammalian-derived Human CD274 proteins Met1-Thr239, with an C-terminal His
<b>Calculated MW</b>	26.1 kDa
<b>Observed MW</b>	35-40 kDa
<b>Accession</b>	Q9NZQ7
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human CD274 proteins, 2 µg/lane of Recombinant Human CD274 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 26.1 kDa.

### Background

Programmed cell death ligand 1(CD274,or B7-H1,PD-L1),is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with an their CD28 family receptors,the B7s are key regulators of the adaptive immune response. CD274 is suggested a negative regulator of T and B cell,and play important role in mediating tolerance of lymphocytes to self-antigens. It also involved in the costimulatory signal,essential for T-cell proliferation and production of IL10 and IFNG,in an IL2-dependent and a PDCD1-independent manner. PD-L1 is a 290 aa transmembrane protein with an a calculated molecular weight of 33 kDa,the apparent molecular weight has been reported as 45-70 kDa,suggesting probable glycosylation .

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017