AF/LE Purified Anti-Human CD48 Antibody[156-4H9]


catalog number: E-AB-F10610



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD48 protein |
| Host | Mouse |
| Isotype | Mouse IgG2a, κ |
| Clone | 156-4H9 |
| Purification | >98%, Protein A/G purified |
| Conjugation | None (AF/LE) |
| buffer | Sterile PBS, pH 7.2. $<$ 1.0 EU per mg of the antibody as determined by the LAL method |

| Applications | Recommended Dilution |
|--------------|---|
| FCM | $2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ |
| Data | |



Human peripheral blood lymphocytes were stained with 0.2 μ g AF/LE Purified Anti-Human CD48 Antibody[156-4H9] (Right) and 0.2 μ g Mouse IgG2a, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

IgG Secondary Antibody.

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
| | thaw cycles. This preparation contains no preservatives, thus it should be handled |
| | under aseptic conditions. |
| Shipping | Order now, ship in 3 days |
| Background | |

This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only