

Biotin Anti-Human CD40 Antibody[G28.5]

Catalog Number: E-AB-F1214B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	G28.5
Isotype Control	Biotin Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD19 Antibody and Biotin Anti-Human CD40 Antibody followed by Streptavidin-Elab Fluor® 647 (Left). Lymphocytes are stained with FITC Anti-Human CD19 Antibody and Biotin Mouse IgG1, κ Isotype Control followed by Streptavidin-Elab Fluor® 647 (Right).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Bp50;CD40L receptor;CDw40;TNFRSF5
Uniprot ID	P25942
Gene ID	958

For Research Use Only

Background

CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand).