

Recombinant Cyclin D2 Monoclonal Antibody

catalog number: **AN301830L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Rat;
Immunogen	Recombinant human Cyclin D2 fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A542
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution


WB	1:2000-1:10000
IHC	1:100-1:1000

Data




Western Blot with Cyclin D2 Monoclonal Antibody at dilution of 1:10000. Lane 1: HEK-293, Lane 2: HeLa, Lane 3: Caco-2


Observed-MW:36 kDa
Calculated-MW:33 kDa



Immunohistochemistry of paraffin-embedded Human colon cancer using Cyclin D2 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Human ovarian cancer using Cyclin D2 Monoclonal Antibody at dilution of 1:100.



Immunohistochemistry of paraffin-embedded Rat stomach using Cyclin D2 Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Cyclin D2 (CCND2) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. CCND2 forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. CCND2 has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. High level expression of CCND2 was observed in ovarian and testicular tumors.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1