Elabscience®

CD164 Monoclonal Antibody

catalog number: AN200126P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human CD164 Protein
Host	Mouse
Isotype	IgGl
Clone	10A4
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS
Applications	Recommended Dilution
IHC-P	1:50-1:200
FCM	1:25-1:100

Data



Immunohistochemistry of paraffin-embedded human prostatic carcinoma using CD164 Monoclonal Antibody at dilution of 1:60.

Flow cytometric analysis of Human CD164 expression on human whole blood monocytes. Cells were stained with purified anti-Human CD164, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter



Immunohistochemistry of paraffin-embedded human pancreatic carcinoma using CD164 Monoclonal Antibody at 1.1 .. C 1 CO

dilution of	of 1:60.		
Preparation & Storage			
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.		
Shipping	Ice bag		
Background			
For Research Use Only			
Toll-free: 1-888-852-8623	Tel: 1-832-243-6086	Fax: 1-832-243-6017	

Web:www.elabscience.com

Email:techsupport@elabscience.com

Elabscience®

Sialomucin core protein 24 also known as endolyn or CD164 (cluster of differentiation 164) is a novel 80- to 90-kD mucinlike molecule expressed by human CD34+ hematopoietic progenitor cells. Isoform 1 and isoform 3 of CD164 are expressed in hematopoietic and non-hematopoietic tissues. Isoform 1 is expressed by prostate cancer tumors and prostate cancer cell lines. The expression is greater in bone metastases than in primary tumors. Expression in osseous metastasis is greater than that in soft tissue metastasis. Isoform 2 of CD164 is expressed in the small intestine, colon, lung, thyroid and colorectal and pancreatic adenocarcinoma. Isoform 4 is expressed by both hematopoietic progenitor cells and bone marrow stromal cells. CD164 belongs to the CD164 family. The cluster of differentiation (cluster of designation) (often abbreviated as CD) is a protocol used for the identification and investigation of cell surface molecules present on white blood cells initially but found in almost any kind of cell of the body, providing targets for immunophenotyping of cells. CD164 may play an important role in prostate cancer metastasis and the infiltration of bone marrow by cancer cells. CD164 promotes myogenesis by enhancing CXCR4-dependent cell motility. This protein positively regulates myoblast migration and promotes myoblast fusion into myotubes. CD164 may play a key role in hematopoiesis by facilitating the adhesion of CD34+ cells to bone marrow stroma and by negatively regulating CD34+ hematopoietic progenitor cell growth.