

Recombinant Human Thymic Stromal Lymphopoietin/TSLP (R127A, R130A, C-10His)

Catalog Number: PKSH033961

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Thymic Stromal Lymphopoietin;TSLP protein Tyr29-Gln159(R127A, R130A), with an C-terminal His
Calculated MW	16.1 kDa
Observed MW	20-27 kDa
Accession	Q969D9
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Histidine-HCl, 4% Sucrose, 4% Mannitol, 0.02% Tween 80 (w/v), pH6.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Thymic stromal lymphopoietin (TSLP) is a novel member of the hemopoietic cytokine family that promotes the development of B cells and shares overlapping activity with IL-7. The human TSLP protein comprises a 28 amino acids (aa) signal sequence and 131 aa mature region. Human TSLP has two isoforms lftTSLP and sftTSLP produced by alternative splicing. lftTSLP is expressed in a number of tissues including heart, liver and prostate, and sftTSLP (63aa) is predominantly expressed in keratinocytes of oral mucosa, skin and in salivary glands. In aa sequence level, Human TSLP displays about 43% identity with mouse TSLP. TSLP is a cytokine that functions mainly on myeloid cells; it induces the release of T cell-attracting chemokines from monocytes and enhances the maturation of CD11c(+) dendritic cells. TSLP has proliferative effects on the myeloid cell line and may initiate asthma or atopic dermatitis responses by directly activating mast cells. TSLP signals cells via the interleukin-7 receptor- α chain (IL-7R α), shared with IL-7, together with the TSLP receptor (TSLPR) subunit. Recent studies indicate that TSLP and its receptor are novel therapeutic targets for rheumatoid arthritis, for increased intraarticular TSLP concentrations in patients has caused chemotaxis and activation of arthritogenic T cells.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017