A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219]

Catalog Number: E-AB-F1093Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

lsotype Rat lgG2a, κ

Clone No. R17 217.1.3/TIB-219

Isotype Control Elab Fluor® Violet 450 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832Q]

Conjugation Elab Fluor® Violet 450

Conjugation Information Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



C57BL/6 mouse bone marrow cells are stained with APC

Anti-Mouse TER-119 Antibody and Elab Fluor® Violet 450 Anti-Mouse CD71 Antibody (Left). Splenocytes are stained

with APC Anti-Mouse TER-119 Antibody and Elab Fluor® Violet 450 Rat IgG2b Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD71;TR;TfR1;TfR1;Tfrc;Transferrin receptor protein 1;Trfr

 Uniprot ID
 Q62351

 Gene ID
 22042

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD71 is a 95 kD type II heterodimeric transmembrane glycoprotein that is also known as T9 and transferrin receptor. CD71 is expressed on proliferating cells, reticulocytes, and erythroid precursors. Its expression is very low on resting leukocytes. CD71 plays a role in the control of cellular proliferation by facilitating the uptake of iron via ferrotransferrin binding and the recycling of apotransferrin to the cell surface.

Fax: 1-832-243-6017