## **Elabscience**®

### **MRPS18B** Polyclonal Antibody


#### catalog number: E-AB-18806

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Fusion protein of human MRPS18B
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

WB	1:500-1:2000
IHC	1:25-1:100

Data





Western blot analysis of HepG2 Jurkat and 231 cell lysates using MRPS18B Polyclonal Antibody at dilution of 1:200

**Observed-MW:Refer to figures** 

Immunohistochemistry of paraffin-embedded Human tonsil tissue using MRPS18B Polyclonal Antibody at dilution of 1:55(×200)

# Calculated-MW:29 kDa Preparation & Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Shipping The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

#### Background

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrian. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that belongs to the ribosomal protein S18P family. The encoded protein is one of three that has significant sequence similarity to bacterial S18 proteins. The primary sequences of the three human mitochondrial S18 proteins are no more closely related to each other than they are to the prokaryotic S18 proteins. Pseudogenes corresponding to this gene are found on chromosomes 1q and 2q.