Recombinant Human BLNK/Ly-57 Protein (His Tag)

Catalog Number: PKSH033753

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Human		
Source	E.coli-derived Human BLNK; Ly-57 protein Met1-Ser456, with an C-terminal His		
Calculated MW	51.5 kDa		
Observed MW	40-80 kDa		
Accession	AAH18906		
Bio-activity	Not validated for activity		
Properties			
Purity	> 90 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Data

kDa	МК	R
120	-	
90	THE OWNER	
60	- 25.00	Contraction of the local division of the loc
40	-	A CONTRACTOR
40	6426	
30	-	and a state of the state of the
		and the second
20	and it	and a state of the
20	1000	12 2 3 3 3
14	1 Sector	100000000
14		ALCONTRACTOR OF
	Colores .	
	1.00	
		Carton Start

> 90 % as determined by reducing SDS-PAGE.

Background

B-Cell Linker Protein (BLNK) is a cell membrane protein which contains 1 SH2 domain. BLNK is expressed in B cells and fibroblast cell lines, playing a important role in B cell receptor signaling. BLNK as a central linker protein, downstream of the B-cell receptor (BCR), bridges the SYK kinase to a multitude of signaling pathways and regulating biological outcomes of B-cell function and development. BLNK associates with the activation of ERK/EPHB2, MAP kinase p38 and JNK, modulates AP1, NF-kappa-B and NFAT activation. BLNK involves in BCR-mediated PLCG1 and PLCG2 activation and Ca2+ mobilization and is required for trafficking of the BCR to late endosomes. BLNK deficiency results in agammaglobulinemia type 4 and much more profound block in B-cell development.

For Research Use Only Toll-free: 1-888-852-8623