

Recombinant Rhesus macaque CD27/TNFRSF7 Protein (Fc Tag)

Catalog Number: PKSQ050077

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rhesus macaque
Source	HEK293 Cells-derived Rhesus macaque CD27/TNFRSF7 protein Thr21-Ile192, with an C-terminal Fc
Calculated MW	46.5 kDa
Observed MW	60-90 kDa
Accession	F7BYS2
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

CD27 antigen, also known as CD27L receptor, T-cell activation antigen CD27, T14, S152, Tp55, TNFRSF7 and Tumor necrosis factor receptor for superfamily member 7, belongs to the TNF-receptor superfamily. CD27 is a single-pass type I membrane protein and exists as a homodimer form, containing three TNFR-Cys repeats. CD27 transduces signals that lead to the activation of NF-KappaB and MAPK8/JNK. CD27 is involved in regulating B-cell activation and immunoglobulin synthesis, binding to the ligand CD70. TRAF2 and TRAF5 have been shown to mediate the signaling process of CD27. CD27-binding protein (SIVA), which is a proapoptotic protein, can bind to CD27 and is thought to play a key role in the apoptosis. CD27 is required for generation and long-term maintenance of T cell immunity.

For Research Use Only