

## FITC Anti-Mouse CD6 Antibody[OX-129]

Catalog Number: AN00327C

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, κ
<b>Clone No.</b>	OX-129
<b>Isotype Control</b>	FITC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832C]
<b>Conjugation</b>	FITC
<b>Conjugation Information</b>	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of C57BL/6 murine splenocytes cells with Elab Fluor® Violet 450 Anti-Mouse CD3 Antibody and FITC Anti-Mouse CD6 Antibody[OX-129] (left) or FITC Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	CD6/TP120
<b>Uniprot ID</b>	Q61003
<b>Gene ID</b>	12511

### For Research Use Only

## Background

CD6, a 100 kD monomeric T cell surface glycoprotein, is a member of the scavenger receptor cysteine-rich protein superfamily. It is expressed on mouse thymocytes and splenic T cells and neurons, but not on splenic B cells. CD6 has a long cytoplasmic tail in mice, with two proline-rich domains that interact with the -SH3 domain binding sequence. CD6 binding to its ligand, CD166 (ALCAM), serves as a costimulatory molecule through its interaction with SLP-76. CD6 mediated signaling may contribute to thymocyte survival and functional avidity in mice and men.