TRO Polyclonal Antibody

catalog number: E-AB-53006



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|-----------------------|--|
| Description | |
| Reactivity | Human |
| Immunogen | Fusion protein of human TRO |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:50-1:200 |
| Data | |
| 5 | of paraffin-embedded Human liver D Polyclonal Antibody at dilution of |
| T | 1:80(×200) |
| Preparation & Storage | |
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |
| Background | |
| This gene encodes a m | embrane protein that mediates cell adhesion between trophoblastic cells and the epithelial cells of |
| the endometrium. The | encoded protein participates in cell signalling during embryo implantation, and may also be |

the endometrium. The encoded protein participates in cell signalling during embryo implantation, and may also be involved in cancer formation. This gene is located near several other closely related genes on chromosome X. Alternative splicing results in multiple transcript variants.

For Research Use Only