

## Recombinant Rat GITR/TNFRSF18 Protein (Fc Tag)

**Catalog Number:** PKSR030328

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat GITR/TNFRSF18 protein Met1-Lys121, with an C-terminal hFc
<b>Calculated MW</b>	37.7 kDa
<b>Observed MW</b>	47 kDa
<b>Accession</b>	Q5M835
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

GITR, also known as TNFRSF18(CD357), belongs to the tumor necrosis factor receptor (TNF-R) superfamily. It is the receptor for TNFSF18. GITR plays a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. GITR may be involved in interactions between activated T-lymphocytes and endothelial cells and in the regulation of T-cell receptor-mediated cell death. GITR and its ligand are important costimulatory molecules in the pathogenesis of autoimmune diseases. It also mediates NF-kappa-B activation via the TRAF2/NIK pathway.

### For Research Use Only