

A Reliable Research Partner in Life Science and Medicine

# FITC Anti-Mouse CD103 Antibody[M290]

Catalog Number: E-AB-F1090C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ Clone No. M290

FITC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832C] Isotype Control

Conjugation

**Conjugation Information** FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### **Applications** Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD3 Antibody and FITC Anti-Mouse CD103 Antibody (Left). Splenocytes are stained with APC Anti-Mouse CD3 Antibody and FITC Rat IgG2a, κ Isotype Control (Right).

#### **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

#### **Antigen Information**

**Alternate Names** CD103;Integrin alpha M290;Integrin alpha-E;Itgae

**Uniprot ID** Q60677 Gene ID 16407

## For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

### **Background**

CD103 is a type I transmembrane glycoprotein known as  $\alpha E$  integrin or Integrin  $\alpha IEL$  chain. It belongs to the integrin family and is primarily found on intestinal intraepithelial lymphocytes (IEL). CD103 is also expressed on a subpopulation of lamina propria T cells, epithelial dendritic cells, lamina propria-derived dendritic cells, and a small subset of peripheral lymphocytes. T regulatory cells express high level of CD103. The CD103 expression on lymphocytes can be induced upon activation and TGF- $\beta$  stimulation. In association with integrin  $\beta 7$ , CD103 is expressed as  $\alpha E/\beta 7$  heterodimer. Mature CD103 protein can be cleaved into 2 chains, a 150 kD (C-terminal) chain and a 25 kD (N-terminal) chain, which remain linked by disulfide bonds. CD103 binds to E-cadherin and mediates homing of lymphocytes to the intestinal epithelium.