

Recombinant Human LEPTIN Protein(GST Tag)

Catalog Number: PDEH101125

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human LEPTIN protein Val22-Cys 167, with an N-terminal GST
Calculated MW	41.9 kDa
Observed MW	45 kDa
Accession	P41159
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human LEPTIN proteins, 2µg/lane of

Recombinant Human LEPTIN proteins was resolved with
SDS-PAGE under reducing conditions, showing bands at 45
kDa

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1

Leptin is one of the most important hormones secreted by adipocytes, as an adipokine that modulates multiple functions including energy homeostasis, thermoregulation, bone metabolism, endocrine, and pro-inflammatory immune responses. The circulating leptin levels serve as a gauge of energy stores, thereby directing the regulation of energy homeostasis, neuroendocrine function, and metabolism. Recent studies suggest that leptin is physiologically more important as an indicator of energy deficiency, rather than energy excess, and may mediate adaptation by driving increased food intake and directing neuroendocrine function to conserve energy, such as inducing hypothalamic hypogonadism to prevent fertilization. One of these functions is the connection between nutritional status and immune competence. The adipocyte-derived hormone Leptin has been shown to regulate the immune response, innate, and adaptive response, both in normal and pathological conditions. Thus, Leptin is a mediator of the inflammatory response. Leptin has a dual effect on bone, acting by two independent mechanisms. As a signal molecule with growth factor characteristics, leptin can stimulate osteoblastic cells and inhibit osteoclast formation and activity, thus promoting osteogenesis. However, as a molecule that stimulates sympathetic neurons in the hypothalamus, leptin indirectly inhibits bone formation. This inhibitory effect of leptin mediated by activation of the sympathetic nervous system can be abrogated by the application of blood pressure-reducing beta-blockers, which also inhibit receptors of hypothalamic adrenergic neurons. Leptin appears to regulate some features defining Alzheimer's disease (AD) at the molecular and physiological level. Leptin can stimulate mitogenic and angiogenic processes in peripheral organs. Because leptin levels are elevated in obese individuals and excess body weight has been shown to increase breast cancer risk in postmenopausal women. Furthermore, a recent report clearly shows that targeting leptin signaling may reduce mammary carcinogenesis.