Recombinant Human NCR3/NKp30 Protein (Fc Tag)

Catalog Number: PKSH032787



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 1	00	cri	m	17	٦m
J			174	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 40.2 kDa

 Accession
 O14931

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Natural Cytotoxicity Triggering Receptor 3 (NCR3) along with NKp44 and NKp46 constitute a group of receptors termed "Natural Cytotoxicity Receptors". They play a major role in triggering NK-mediated killing of most tumor cells lines. NKp30 is a type I transmembrane protein having a single extracellular V-like immunoglobulin domain. NKp30 is selectively expressed both in resting and activated human NK cells. In addition; NKp30 is also involved in NK-mediated induction of dendritic cell (DC) maturation. It has been demonstrated that NK cell activation signaling specifically induces lytic activity against several tumor cell types and synthesis of new NF-kB dependent proteins during the initiation of cytotoxicity.

For Research Use Only