Purified Anti-Human CD52 Antibody[Campath-1G]

catalog number: AN004360P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description					
Reactivity	Human				
Immunogen	Recombinant Human CD52 protein				
Host	Rabbit				
Isotype	Rabbit IgG, κ				
Clone	Campath-1G				
Purification	>98%, Protein A/G purified				
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze				
	to completely remove the stabilizer prior to labeling.				
American					

Applications	Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data

	429							
	300					Λ		
Count	8							
	8					\mathbb{N}		
	0	L.,	1		- And	1		
	10	01	102	103	104	105	106	107
			CE	52 Ale	xa Fluc	r® 647	н	

Human peripheral blood lymphocytes were stained with 0.2

μg Purified Anti-Human CD52 Antibody[Campath-1G]

(Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left),

followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

IgG Secondary Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	Ice bag
Desleanend	

Background

CD52, also known as Cambridge pathology antigen 1 (CAMPATH-1), is a 25-29 kD glycoprotein containing a large Nlinked carbohydrate moiety. The actual molecule of CD52 is only 8-9 kD. It is expressed in the male reproductive tract and on virtually all lymphocytes (T and B cells), as well as macrophages/monocytes, eosinophils, and red cells. CD52 is thought to play a role in carrying and orienting carbohydrates. CD52 is a potent target for complement-mediated lysis and antibody-mediated cellular cytotoxicity and has been used as a depletion target for chronic lymphocytic leukemia (CLL)/lymphoma and immunosuppression.

For Research Use Only