

TTC12 Polyclonal Antibody

catalog number: E-AB-52773

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Fusion protein of human TTC12
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution

WB	1:500-1:2000
IHC	1:30-1:150


Data



Western blot analysis of Human bladder transitional cell carcinoma grade 2-3 tissue lysate using TTC12 Polyclonal Antibody at dilution of 1:300

Observed-MV: Refer to figures

Calculated-MV: 79 kDa



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TTC12 Polyclonal Antibody at dilution of 1:40 (x200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

TTC12 (tetratricopeptide repeat domain 12), also known as TPARM, is a 705 amino acid cytoplasmic protein containing a TPR repeat domain and three armadillo repeat motifs. TTC12 is expressed in testis, prostate, lung, germinal center B cells, neuroblastoma, teratocarcinoma, colon cancer and gastric cancer. The gene encoding TTC12 is located in a region of human chromosome 11 that is commonly deleted in a variety of cancers, therefore, it is considered a candidate tumor suppressor gene. Human chromosome 11 houses over 1,400 genes and comprises nearly 4% of the human genome.

For Research Use Only