

HDAC3 Polyclonal Antibody

catalog number: **E-AB-64237**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human HDAC3 (NP_003874.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200
IF	1:50-1:200


Data





Western blot analysis of extracts of various cell lines using HDAC3 Polyclonal Antibody at dilution of 1:1000.

Observed-MW:48 kDa

Calculated-MW:48 kDa/49 kDa




Immunohistochemistry of paraffin-embedded Mouse liver using HDAC3 Polyclonal Antibody at dilution of 1:100 (40x lens).




Western blot analysis of extracts of Mouse brain using HDAC3 Polyclonal Antibody at dilution of 1:1000.

Observed-MW:48 kDa

Calculated-MW:48 kDa/49 kDa




Immunohistochemistry of paraffin-embedded Human lung cancer using HDAC3 Polyclonal Antibody at dilution of 1:100 (40x lens).



For Research Use Only

Immunohistochemistry of paraffin-embedded Human colon carcinoma using HDAC3 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human stomach using HDAC3 Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunofluorescence analysis of HeLa cells using HDAC3 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the histone deacetylase/acuc/alpha family. It has histone deacetylase activity and represses transcription when tethered to a promoter. It may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. This protein can also down-regulate p53 function and thus modulate cell growth and apoptosis. This gene is regarded as a potential tumor suppressor gene.

For Research Use Only