A Reliable Research Partner in Life Science and Medicine

# Recombinant Rhesus macaque IL-33/Interleukin-33 Protein (His Tag)

Catalog Number: PKSQ050072

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$					٠.			
H))	es	C	r٦	n	tı	n	m	

**Species** Rhesus macaque

Source E.coli-derived Rhesus macaque IL-33/Interleukin-33 protein Ser112-Ile270, with an N-

terminal His

Mol\_Mass 20.3 kDa Accession F7DLG4

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



> 95 % as determined by reducing SDS-PAGE.

## **Background**

Interleukin-33 (IL-33) belongs to the IL-1 superfamily. IL33 is highly expressed in endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes, but almost undetectable in placenta. IL33 induces the production of T helpe r-2 (Th2)-associated cytokines. IL-33 is a cytokine that mediates its biological effects by binding to and signals through IL1RL1/ST2 and IL-1 Receptor Accessory Protein (IL1RAP), activating intracellular molecules in the NF-κB and MAP kinase signaling pathways that drive production of type 2 cytokines from polarized Th2 cells.

## For Research Use Only